

Collection Of Mid-Stream Urine For Bacterial Culture & Sensitivity 中段尿液細菌和藥物測試

Preparation and Instruction for Investigation:

- 1. Patient is advised to wash hands pre-procedure.
- 2. For female patient, hold labia apart and pass the first portion voided urine into toilet and collect midstream urine into a clean polypropylene container without preservatives to 1/2 full.
- 3. For male patient, retract foreskin and pass the first portion voided urine into toilet and collect midstream urine into a clean polypropylene container without preservatives to 1/2 full.
- 4. Close the lid of the container tightly and label appropriately.
- 5. Please send specimen together with request form to the Clinical Laboratory. Specimen should deliver as soon as possible. Please store at 4 8°C if not deliver immediately.

測試準備及須知:

- 1. 病人需於採集標本前潔手。
- 2. 女性應拉開外陰部或男性應拉緊包皮將前段尿液排走棄置,將中段尿液留於已消毒的聚丙烯容器至 1/2 瓶滿。
- 3. 將容器蓋好及扭緊, 確定樣本不會滲漏。
- 4. 在該容器適當位置貼上病人標籤。
- 5. 請盡快把樣本連同檢驗單送往化驗室。若無法立刻送往化驗室,請保存樣本 4-8℃。